

# The LEGEND Neutrinoless Double Beta Decay Experiments

LEGEND




Large Enriched  
Germanium Experiment  
for Neutrinoless  $\beta\beta$  Decay

Chris Jillings

2024-07-31

SNOLAB SEF/EAC meeting





CIEMAT  
 Comenius Univ.  
 Czech Tech. Univ. Prague and IEAP  
 Daresbury Lab.  
 Duke Univ. and TUNL  
 Gran Sasso Science Inst.  
 Indiana Univ. Bloomington  
 Inst. Nucl. Res. Rus. Acad. Sci.  
 Jagiellonian Univ.  
 Joint Inst. for Nucl. Res.  
 Joint Res. Centre Geel  
 Lab. Naz. Gran Sasso  
 Lancaster Univ.  
 Leibniz Inst. for Crystal Growth

Leibniz Inst. for Polymer Research  
 Los Alamos Natl. Lab.  
 Max Planck Inst. for Nucl. Phy.  
 Max Planck Inst. for Physics  
 Natl. Res. Center Kurchatov Inst.  
 Natl. Res. Nucl. Univ. MEPHI  
 North Carolina State Univ.  
 Oak Ridge Natl. Lab.  
 Polytech. Univ. of Milan  
 Princeton Univ.  
 Queen's Univ.  
 Roma Tre Univ. and INFN  
 Simon Fraser Univ.  
 SNOLAB

South Dakota Mines  
 Tech. Univ. Dresden  
 Tech. Univ. Munich  
 Tennessee Tech. Univ.  
 Univ. of California and LBNL  
 Univ. College London  
 Univ. of L'Aquila and INFN  
 Univ. of Cagliari and INFN  
 Univ. of California San Diego  
 Univ. of Houston  
 Univ. of Liverpool  
 Univ. of Milan and INFN  
 Univ. of Milano Bicocca and INFN  
 Univ. of New Mexico

Univ. of North Carolina at Chapel Hill  
 Univ. of Padova and INFN  
 Univ. of Regina  
 Univ. of South Carolina  
 Univ. of South Dakota  
 Univ. of Tennessee  
 Univ. of Texas at Austin  
 Univ. of Tuebingen  
 Univ. of Warwick  
 Univ. of Washington and CENPA  
 Univ. of Zurich  
 Williams College


# Outline

- Double Beta Decay with Germanium Detectors
- LEGEND Concept and Background Control
- LEGEND-200 Results and Plans
- LEGEND-1000 Development

$0\nu\beta\beta$  candidates are even-even nuclei as the mass parabola for odd-odd nuclei is shifted

76 AMU

Atomic Mass for A=76




Odd odd nuclei

Even even nuclei

Single beta decay to As is energetically forbidden

75.9 AMU

Ge crystals with point-contact and electronics near crystal allows for exceptional pulse shape discrimination while maintaining energy resolution.




Combined energy calibration for DS0-6


**Majorana Demonstrator**

<https://arxiv.org/abs/1902.02299>

Zoom into TI-208 peak




# Ge-76 has a long history in $0\nu\beta\beta$ searches



PRL 130, 062501 (2023)  
 $T_{1/2} > 8.3 \times 10^{25}$  yr (90%CL)


Data tabulated in review by Avignone and Elliott.  
<https://doi.org/10.3389/fphy.2019.00006>  
<https://arXiv.org/abs/1901.02805>

LEGEND uses sophisticated large enriched Ge-76 detectors building on work by Majorana and GERDA.




# Many techniques are used to control background:

- Bare crystals with small-mass electronics (ASIC) near crystal: Exquisite energy resolution
- PSD in analysis to reject multi-site events
- Crystals in instrumented liquid-argon bath for cooling and Compton rejection. (Atmospheric Argon for LEGEND-200 and Underground Argon for LEGEND-1000)
- Instrumented atmospheric argon shield
- Instrumented water shielding tank






# LEGEND-1000 Background Model




# A sample 10 t-yr synthetic data set illustrates discovery potential




Discovery: a 50% chance or greater that a 10 tonne-year results in a signal  $3\sigma$  above null hypothesis

LEGEND-1000 is designed to have  $0\nu\beta\beta$  discovery potential at a  $10^{28}$  year half life



# Strategy for Suppressing Ge-77m Background (from cosmogenic activation of Ge-76)

- Acrylic panels are added to design in outer argon detector to thermalize neutrons
- A method for tagging cosmogenic fast neutrons has been developed in the instrumented outer (atmospheric) argon.
- Only one branch of Ge-77m produces a single-site event in LEGEND. Use a veto after a neutron signal.
- With a loss of  $\sim 3\%$  detection efficiency, the total background rate at LNGS similar to that at SNOLAB.
- With this improvement in place, the physics reach at LNGS and SNOLAB are very similar.
- The detailed study will be published shortly
- A search for Ge-77m in LEGEND-200 is underway – will be upcoming publication.



# LEGEND-200 commissioning showing crystals and liquid argon readout




E. Sacchetti




Photo: E. Sacchetti

**60 kg campaign:**  
 First operation of 60 kg of HPGe detectors and full LAr instr.  
 Final hardware optimisations  
 Special calibration runs



**142 kg installation:**  
 Installation of all available HPGe detectors as well as full LAr installation, DAQ, readout electronics





**LAr instrumentation:**  
 Construction & commissioning of LAr instr. hardware & readout electronics.




# Data Release at Neutrino 2024

## THE LEGEND-200 EXPERIMENT AT LNGS



### Hardware status — see talk at [TAUP23]

- Installed first **142 kg** of HPGe detectors (Oct '22)
- 130 kg operational (12 kg OFF due to hardware issues)




The first year of LEGEND-200 physics data in the quest for  $0\nu\beta\beta$  • L. Pertoldi • Neutrino 2024, Milano • 18 June 2024

# Results Presented at Neutrino 2024

MODELING DATA BEFORE ANALYSIS CUTS [SILVER]

POSTER THE LEGEND-200 BACKGROUND MODEL • T. Dixon, S. Calgaro




- Simulations and material radioassay **underpredict**  $^{228}\text{Th}$  in physics data
  - *Hard to estimate systematic uncertainty on the assay results*
  - *ICP-MS not predictive if secular equilibrium is broken*
- This background is efficiently **suppressed by analysis cuts**

# Results Presented at Neutrino 2024

MODELING DATA BEFORE ANALYSIS CUTS [SILVER]


POSTER THE LEGEND-200 BACKGROUND MODEL • T. Dixon, S. Calgario



- Bayesian background model using data before analysis cuts [SILVER]
  - Includes 10.2 kg yr from special “background characterization” runs
- Data well reproduced, model is flat at  $Q_{\beta\beta}$ 
  - No “hotspot” or significant asymmetry observed in data
  - Model can test hypotheses on the origin of  $^{228}\text{Th}$




## DATA AFTER PULSE SHAPE DISCRIMINATION [GOLDEN]




- Strong **suppression of surface  $\alpha$  and  $\beta$  ( $^{42}\text{K}$ ) events**
- ~60% suppression of Compton multi-site events at  $Q_{\beta\beta}$
- $0\nu\beta\beta$  survival fraction of ~85%

## DATA AFTER PULSE SHAPE DISCRIMINATION AND ARGON ANTI-COINCIDENCE CUT [GOLDEN]



LEGEND-200 · 06-2024

- Strong **anti-correlation** of argon and PSD cuts
- Overall  $0\nu\beta\beta$  survival fraction of  $\sim 60\%$
- **“Pure”  $2\nu\beta\beta$  distribution, few events surviving at  $Q_{\beta\beta}$**




LEGEND-200 · 06-2024

# Results Presented at Neutrino 2024

## DATA IN THE REGION OF INTEREST!



5 events surviving in the  
“background estimation window”



# Results Presented at Neutrino 2024



**DATA IN THE REGION OF INTEREST — AFTER UNBLINDING LAST WEEK!**

- 7 events surviving. Background index  
 $BI = 5.3 \pm 2.2 \cdot 10^{-4}$  cts / (keV kg yr)

**PRELIMINARY!**


## GERDA, MAJORANA and LEGEND combined fit

- $p$ -value of background-only = 26%
- $T_{1/2}^{0\nu}$  lower limits (90% frequentist C.L.)

Observed	Sensitivity
$> 1.9 \cdot 10^{26}$ yr	$2.8 \cdot 10^{26}$ yr

## LEGEND-200 contribution

- +30% of limit median expectation
- event at  $1.4 \sigma$  from  $Q_{\beta\beta}$  weakens combined limit



- The DOE launched an ongoing process to examine alternatives for LEGEND-1000.
- Draft recommendations include
  - LEGEND-1000 is needed to meet the  $10^{28}$  year half life sensitivity and therefore cover the inverted hierarchy space. (LEGEND-200 will reach  $10^{27}$  years.)
  - LNGS is the baseline location. The LNGS site has reduced cost to the DOE while maintaining the physics goal.

The mid-scale proposal to the NSF is proceeding well. A new work breakdown structure was created to define responsibilities to be funded under the NSF. A full proposal to NSF was invited and efforts are proceeding accordingly. The site visit and reverse site visit have already taken place and we expect to hear results in October.

The Department of Energy CD-1 review was pushed to November 2024 (from its scheduled date in June 2024). Since the January 2024 EAC meeting there has been a Directors' review and a "red team" review of the CD-1 presentations. These were successful and useful. However, the documentation required to fulfill CD-1 requirements could not be met in parallel with the NSF activity, resulting in the delay to November.

The work on LEGEND at SNOLAB has been scaled back, given the emphasis on deployment at LNGS. Therefore, the efforts will be focused on low-background work: assays and some studies that are consistent with the broader SNOLAB program. The fraction of effort to be spent on LEGEND by Jillings will shrink.

- LEGEND-200 is running at LNGS with recent results presented at Neutrino 2024.
- LEGEND-1000 is in advanced design for deployment at LNGS with SNOLAB as the alternate site.
- 1000 kg of enriched Ge crystals with exquisite energy resolution in an ultra-low background environment have discovery potential with a half life of  $10^{28}$  years for  $0\nu\beta\beta$  in  $^{76}\text{Ge}$ .