

Event by Event classification of alpha-n and IBD Interactions at SNO+

In the study of reactor and geo antineutrinos, tagging of the inverse beta decay (IBD) positron-neutron coincidence signature allows for the elimination of most backgrounds. In many detectors, the primary remaining background is caused by captures on ^{13}C —so called $(\bar{\nu}, n)$ events —which release a neutron and closely mimic the IBD's signature. The most common $(\bar{\nu}, n)$ prompt event is produced by protons recoiling from the neutron, which gives rise to a distinct pulse shape compared to that of the positron from an IBD. A powerful classifier is thus presented, able to purify the IBD signal from most of its $(\bar{\nu}, n)$ background, by discriminating between these pulse shapes. Particular attention is paid to the construction of appropriate training data from Monte-Carlo simulations. The tuning of the β and proton scintillation timing models in these simulations for SNO+ is also discussed. Tuning of the former is achieved via the selection of a high purity sample of in-situ ^{214}Bi to ^{214}Po decays. The latter makes use of the deployment of a radioactive Americium-Beryllium source. Finally, results of this classification on expected reactor and geo-neutrino signals are shown, and the question of over-tuning is tackled.

Primary author: PAGE, James (U Sussex)

Presenter: PAGE, James (U Sussex)